


BLSC Invasive species and Biohazard prevention policy and Code of Practice.

Author: Canice Keane


Date: July 2017

Version: 1.0



Contents

1.0 Background	3
2.0 Blessington Sailing Club Policy	4
3.0 How can a Code of Practice reduce risk?	4
4.0 Scope of the Code of Practice (Kelly & Maguire, 2008).	4
4.01 Invasive alien species prevention and surveillance.	4
4.02 Inspect – remove – dispose – report.	5
4.03 Small crafts including anglers and divers using boats.	5
4.04 Leisure and industrial craft sectors	6
Bibliography	8
Annex 1: Some high impact invasive alien species (Kelly & Maguire, 2008)	9
Most Unwanted: Didemnum vexillum Didemnum vexillum	9
Most Unwanted: Slipper limpet	11
Most Unwanted: Asian rapa whelk	12
Most Unwanted: Zebra mussel	13
Most Unwanted: Curly leaved waterweed	15
Most Unwanted: Brown rat	17
Most Unwanted: Ship/black rat	18



1.0 Background

Invasive non-native plant and animal species are the second greatest threat to biodiversity worldwide after habitat destruction. They can negatively impact on native species, can transform habitats and threaten whole ecosystems causing serious problems to the environment and the economy.

The Invasive Species Ireland project which began in May 2006 aims to reduce the impact and threats from invasive species on the island of Ireland. This is a joint initiative between National Parks and Wildlife Services and the Northern Ireland Environment Agency. Preventing the spread and new introductions of invasive species is an issue of shared responsibility and requires the involvement of all relevant government agencies, academia, private and voluntary sectors.

Many non-native species do not become invasive but provide considerable benefits to society and industries such as horticulture, agriculture, forestry and aquaculture. Therefore, management of the problem requires engagement and consultation with commercial interests. The development of this code of practice is aimed at reducing the risk from, and impacts of, invasive species and protecting native biodiversity. It is not seeking to stop the use of non- native species or impede the development industry on the island of Ireland. (Kelly & Maguire, 2008)

A risk assessment carried out by Invasive Species Ireland project identified a number of species that are high risk invasive species to the island of Ireland. A number of these organisms are known to contaminate equipment and vessels or are capable of surviving in water and/or fouled ropes and chains that are moved from one water body to another. Some of these mechanisms are quite obvious to the casual observer whereas others can go unseen and unnoticed until an impact is observed in a new area. (See www.invasivespeciesireland.com for more details).

The Irish and UK Governments are contracting parties to a number of international conventions and agreements which require them to take account of invasive introduced species. The Convention on Biological Diversity (CBD) aims to prevent introductions, control invasive species and develop legislation. Other legislative drivers are European Directives (Birds Directive, Habitats Directive and the Water Framework Directive) and national legislation (Wildlife (Amendment) Act 2000 and the Wildlife (Northern Ireland) Order 1985). Both countries are also committed to the EU target of halting the loss of biodiversity by 2010.


Marine invasive species of concern include *Didemnum* species* (a species of invasive tunicate), wireweed* (*Sargassum muticum*), clubbed tunicate* (*Styela clava*), the vase tunicate# (*Ciona intestinalis*), wakame# (*Undaria pinnatifida*), slipper limpet# (*Crepidula fornicata*), and the Asian rapa whelk# (*Rapana venosa*).

Freshwater invasive species of concern include the zebra mussel (*Dreissena polymorpha*)*, floating pennywort (*Hydrocotyle ranunculoides*)*, curly leafed waterweed (*Lagarosiphon major*)*, Nuttalls pondweed (*Elodea nuttallii*)*, species of non-native crayfish#, Asian clam (*Corbicula fluminea*)#, Topmouth Gudgeon (*Pseudorasbora parva*)#, Quagga mussel (*Dreissena bugensis*)#, and the fishhook waterflea (*Cercopagis pengoi*)#.

Terrestrial invasive species of concern that are associated with certain activities of this sector include the ship rat (*Rattus rattus*)* and the brown rat (*Rattus norvegicus*)*. These species can stow away on boats and spread to environmentally sensitive off shore or lake islands where they can impact on nesting birds and biodiversity in general.

* Species known to occur in Ireland

Species currently not known to be present in Ireland
(Kelly & Maguire, 2008)



2.0 Blessington Sailing Club Policy

It is the Policy of the Blessington Sailing Club to implement and adhere to the Invasive species Ireland Water Users Code of Practice in all aspects relating to the purpose of Dingy Sailing and Powerboat Rescue craft.

3.0 How can a Code of Practice reduce risk?

The ability of species contaminate equipment such as trailers, canoes, diving gear, sail boats, powerboats and jet skis has and can continue to deliver invasive species to new areas where they can impact on the biodiversity and economy of the region. This sector has a vital role to play in preventing this movement of species through putting in place mechanisms such as;

- a) preventing fouling of vessels,
- b) removing fouling from vessels in a responsible manner where it cannot return to the aquatic environment,
- c) ensuring that all water is drained from the vessel/craft before transferring to another water body,
- d) preventing fouling of ropes and chains by drying them on a regular basis,
- e) not disposing of live material in the aquatic environment, appropriate use of rat guards when docked
- f) ensuring that equipment is not contaminated prior to entering a new water body.

The impact invasive species exert on native species, endangered species and on the conservation goals of designated areas are costly to control and mitigate against. Good practice has the potential to deliver significant benefits in terms of preventing introduction and spread of non-native species, where these are likely to cause problems.

Compliance with this code of practice will help prevent the spread of invasive species and ensure compliance with legislation. Increased awareness will also contribute to surveillance for potential invaders and better information on the distribution of species already established in Ireland. (Kelly & Maguire, 2008)


4.0 Scope of the Code of Practice (Kelly & Maguire, 2008).

The code of practice applies to all Blessington Sailing Club members or guests wishing to use the Blessington Lakes for the purpose of Sailing or Powerboat Rescue.

4.01 Invasive alien species prevention and surveillance.

Water users have a crucial role to play in the prevention and surveillance of invasive alien species;

- a) Do not introduce non-native species to the island of Ireland.
- b) Keep up-to-date on invasive alien species, their impact and distribution in Ireland.
- c) Keep ID guides on our most unwanted species. ID cards and information leaflets will be placed on the Invasive Species Ireland website. You are encouraged to download these, print copies and keep them with you. Use ID guides when you are out and about and cleaning equipment to help identify if an invasive species is present.
- d) Adopt practices that minimize the opportunities of spreading invasive species.



4.02 Inspect – remove – dispose – report.

Removing build-up of plants and animals from equipment is effective at preventing the opportunity of colonization by invasive species. To prevent the spread of invasive species ensure you always follow these guidelines:

- **Clean** all parts of equipment that comes into contact with the water.
- **Remove** any visible plant, fish, animal matter and mud.
- **Do not move** fouled equipment from one area to another.
- **Watch out** for hitchhikers on ropes and chains.
- **Report sightings** of our most unwanted species or any organism you suspect may be a high risk species. More information on these species can be found on www.invasivespeciesireland.com. Photographs can be submitted via the website.
- **Talk about invasive species.** Talking about the impact invasive alien species have on the environment will help raise awareness of the issue and also inform others of their role in preventing the spread of high impact species.

4.03 Small crafts including anglers and divers using boats.

This group encompasses all small crafts used in the marine and freshwater environments. It includes personal water crafts, jet skis, canoes/kayaks, small boats, dinghies and other small vessel types that are not routinely kept in the water after their use.

This sector transfers equipment from one water body to another on a regular basis. It is important that these protocols are followed each and every time a boat or craft leaves the water:

Avoid running your engine in aquatic weeds, where possible.

Inspect your craft for signs of plants, animals or grit and dirt.

Do not move fouled equipment from one area to another.

Keep your boat in the water for as short a time as possible. Removing your boat from the water prevents build up of fouling on the hull and reduces thereby reduces the opportunities for invasive species to colonize.

Remove any visible plant, fish, animal matter and mud.


Dispose of any material in a rubbish bin or skip. Do not return this to the aquatic environment. It may encourage the spread of an invasive species.

Clean all parts of equipment that comes into contact with the water. Always assume that your craft is infested with an invasive species and must be treated accordingly. High pressure wash hull, lines, bumpers, motor, trim tabs, trailer, and all nooks and crannies.

Drain and flush any water e.g. bilge, live wells, bait buckets, cooling system and any water that collects in the bottom of your craft.

Watch out for hitchhikers on ropes and chains and that may have become entangled in your engine.

Dry your boat (if possible) for 5 days (3-4 weeks under damp conditions) if it has been in a known infested water way.
Report sightings.



4.04 Leisure and industrial craft sectors

These sectors have a crucial role to play in the prevention and surveillance of invasive alien species. This sector encompasses larger boats, those that are routinely kept in the water and those that are moved from one waterbody to another.

Prevent fouling. Fouling of vessels is known to reduce efficiency, increase drag and increase fuel consumption. Also, the increased weight to mooring lines caused by fouling organisms may hinder the ease end users have of handling them. It is in the interest of owners to keep fouling off vessels and lines and thereby protecting the aquatic environment from harm caused by translocations of invasive species. If possible, keep boats in the water for short periods of time.

Treat your boat with appropriate antifouling techniques that adhere to the craft manufacturers recommendations and prevent build-up of organisms.

Ensure boats submit to yearly removal of fouling

When treating a craft 100% surface cover with the chosen method is essential. Small areas left available for plant or animal growth can give species the opportunity to colonise new areas.

Antifouling paints are designed to present marine organisms with a barrier to prevent settlement. These compounds are sometimes toxic to humans, aquatic organisms and terrestrial species and care should be taken to follow the guidelines stipulated by the manufacturer at all times.

If mooring lines become heavily fouled remove them from the water, dispose of fouling in a dustbin or skip (do not allow fouling to return to the aquatic environment), and allow the ropes to dry out for a period greater than 48 hours.

Guidelines to follow when renewing/applying hull fouling.


Adapted from the Massachusetts Clean Marina Guide,
<http://www.mass.gov/czm/marinas/guide/macleanmarinaguide.htm>

For more information on antifouling methods and products please consult The Green Blue webpage on antifouling (<http://www.thegreenblue.org.uk/youandyourboat/cleaningandantifouling.asp>).

Work indoors: Where practical, conduct vessel maintenance indoors or under temporarily covered areas where the rain cannot cause runoff. Sheet plastic shelters can be used.

Work away from the water: At a minimum, always move boats inland to an approved work area before scraping or power washing the hull. Do not perform hull maintenance activities on the launch ramp area or in the lift well.

Do not clean your boat while it is in the water: Never dive/use divers to remove growth on the bottom of boat hulls. This not only represents a health and safety concern but cleaning of biofouling by this manner also removes anti-fouling paint and associated pollutants.



Ensure you follow manufacturer's guidance or receive training on the safe use of this equipment.

Tarps and filter cloth: Use tarps and/or filter cloth to catch scrapings and other debris produced during maintenance work. Tarps and cloth are inexpensive "low-tech" methods to collect debris before it can be washed into coastal waters by storm water. Filter cloths are better than tarps when boat work is expected to last longer than one day. Should it rain, the water passes through the cloth instead of washing the debris off the tarp.

Maintain clean and tidy work area: Clean up the designated work area after scraping and painting. Leaving areas cluttered and messy will cause spills and allow pollutants to be tracked outside the work area. Painting: Because hull paints contain toxic pollutants, they should be used with care. Consider the following when painting your boat.

Maintain designated work areas: Restrict mixing of paints, solvents, and reducers, as well as the painting itself, to designated areas that are located on a hard surface and isolated from the weather.

Do not spray paint while boat is on the water: Sprayed paint can be difficult to control. Paint can be inadvertently sprayed into the water and expose aquatic life to toxic chemicals.

Use spray enclosure: A spray enclosure is a permanent shed or temporary structure housing a boat during painting. Spray booths confine overspray and prevent drifting onto other boats, land, or water.

When using traditional paint application techniques use brushes and rollers where possible. Traditional applications are known to reduce air emissions. Some manufacturers stipulate the use of rollers or brushes so always consult manufacturer's guidelines.

When spray painting, use efficient high volume, low pressure (HVLP) spray guns: HVLP sprays are one of the most efficient means for applying paint. Use a spray gun that is rated at 65 percent efficient paint transfer or greater. These spray guns direct more paint onto the intended surface and as a result, less paint gets into the air, and fewer chemicals are released.

Always maintain appropriate use and storage of hazardous materials and waste: Make certain that all painting materials are used strictly according to manufacturers' instructions. Keep covers and caps on paints, thinners, and solvents to minimize the release of toxic compounds.

Clean up paint and supplies: Treat paint spills like oil spills. Clean up immediately with absorbent materials, paper, and/or rags. Since liquid paints are classified as hazardous material, dispose of paint brushes and paint properly.

Remove fouling prior to long distance journeys. If travelling to or from Britain, Europe or further afield remember that the hull and mooring lines may have potentially damaging hitchhikers present. We have a responsibility to protect not only our environment but that of the areas we visit. Before undertaking such a trip, remove all hull fouling, renew antifouling treatments if required and dry all mooring lines completely.

Do not allow rats to stow away on your boat. Rats are capable of sometimes going unnoticed on larger boats. Use of rat guards on mooring lines can prevent access. Should you find a rat on your boat do not allow it to escape and gain access to sensitive offshore or lake islands. Humanely dispatch any rat that stows away on your boat.

Do not discharge untreated bilge water. Untreated bilge water should be discharged in a responsible manner. Bilge water will contain toxic substances and also may contain the microscopic organisms or life cycle stages that are harmful to native ecosystems.

Take action, the success of this code will depend on the level of uptake by the various stakeholders. There are specific actions which should be undertaken to reduce the risk from invasive species and raise awareness of the issue, these include:



Auditing your actions to identify if you can reduce the risk of transporting or spreading an invasive alien species.

Displaying information materials on your equipment or in your clubhouse.

Discuss the invasive alien species issue with your peers and how their actions can reduce spread.

Sign up to the Invasive Species Ireland network and keep up to date with developments.

The implementation of the code will be supported by a range of measures which will be provided through the Invasive Species in Ireland project. These will include:

Provision of education and awareness materials such as posters and leaflets.

A dedicated section on the website where information can be found on high risk invasive species in Ireland.

Provision of this information in hard copy upon request.

Provision of training materials on invasive species and training can be provided upon request.

A programme of stakeholder engagement to get uptake of the code by the sector and in local and central Government procurement.

A publicity campaign to raise awareness of the code and its aims.

Bibliography

Kelly, J., & Maguire, C. (2008). Water Users Code of Practice. *Water Users Code of Practice*. NIEA and NPWS as part of Invasive Species Ireland. Retrieved from Invasivespecies: www.invasivespeciesireland.com




Annex 1: Some high impact invasive alien species (Kelly & Maguire, 2008)

Most Unwanted: *Didemnum vexillum* *Didemnum vexillum*

Habitat: Marine

Threat: Fisheries, protected species and marine ecosystem functioning Status: Established



A colonial sea squirt from the *Didemnum* (pronounced die-DEM-num) group has recently been found fouling boats and marina structures in Ireland. This particular species of *Didemnum* is fast-growing and has the potential to out compete and smother a large number of native species. The threats posed are of concern to conservationists, aquaculture managers, and fisheries.


What is it?

Exact identification of this species is very difficult and can only be undertaken by a trained expert. However, in general native species are not found in large colonies and do not form the candlewax-like dripping structures, as seen in the photographs. Please see the *Habitas* website on Sponges of Britain and Ireland for more info on native species.

The invasive form can be described as long, ropey or beard-like hanging from hard surfaces such as docks, aquaculture facilities, and ship hulls, or may be found as extensive mats covering rocky sea beds (pebbles, cobbles, boulders, and rock outcrops).

Impact

Alter marine habitats.



Interfere with fishing, aquaculture, and other coastal and offshore activities.
Increased fouling of manmade structures such as docks, moorings, and boat hulls.
Overgrow other organisms such as seaweed, scallops, mussels, and oysters that depend on the seafloor for habitat.
May impact on underwater archaeological sites such as ship wrecks.
Produce noxious substances that deter most fish and other animals.
The colonies have been found at water depths ranging from intertidal to continental shelf depths of 65m (213 ft).

Where is it from?

Uncertain at present. The species is known from The Netherlands and France in Europe while worldwide the species is now recognized in Asia, Australasia-Pacific, Europe, and North America.

Where is it found in Ireland?

Currently confirmed present in Carlingford Lough, Co. Louth, Malahide Marina, Co. Dublin and Galway Bay, Co. Galway.

How did it get here?

Uncertain at present. Likely vector is fouling of ocean going vessel and/or contamination of aquaculture produce.

How is it spreading in Ireland?

Once the species becomes established in an area, it can spread rapidly by both sexual reproduction and asexually by fragmentation of the colonies. Species is known to grow on mobile organisms such as crabs. Fouling on sea going vessels is a key vector in dispersal of this species. The species may also establish itself with movement of shellfish for the aquaculture industry.

What you can do?

Do not interfere with colonial tunicates. These species can spread more rapidly when a section of the colony becomes detached and carried with currents to new areas. If you can do so safely, photograph the species.

Everyone is urged to thoroughly wash and dry all equipment that was near an infestation of this species. This is essential to avoid contamination of new areas. It is important to do this where the rinse will not return to the marine environment. New infestations may result if you do not undertake this.

Report all sightings at the Alien Watch page of www.invasivespeciesireland.com.



Most Unwanted: Slipper limpet

Crepidula fornicata

Habitat: Marine

Threat: Threatens native ecosystems and species Status: Potential



This is a species of mollusc that was introduced to Europe in the 19th century. The slipper limpet has also been introduced to Ireland accidentally along with imported oyster spat but so far has failed to become established.


What is it?

This species of mollusc inhabits the area around the low water mark and into the shallow subtidal. It is often found attached to shells of mussels and the native oyster. MarLin describes this species as possessing an oval shell, up to 5 cm in length, with a much reduced spire. The large aperture has a shelf, or septum, extending half its length. The shell is smooth with irregular growth lines and white, cream, yellow or pinkish in colour with streaks or blotches of red or brown. Slipper limpets are commonly found in curved chains of up to 12 animals. Large shells are found at the bottom of the chain, with the shells becoming progressively smaller towards the top.

Impact

In shallow bays where the slipper limpet has been introduced in France, it can completely smother the sediment creating beds with several thousand individuals per m². Dense aggregations of slipper limpet trap suspended silt, faeces and pseudofaeces altering the benthic habitat. Where slipper limpet stacks are abundant, few other bivalves can live amongst them. The slipper limpet is a serious threat to oyster beds because of this.

It has also been observed that live maerl thalli, which are a protected species and form an important protected habitat, become covered in slipper limpets and the spaces between the thalli of the bed become clogged with silt; this kills the maerl thalli and dramatically alters associated communities. No management measures have proven effective for this species in this habitat.



Where is it from?

Originally found on the east coast of the Americas between Canada and Mexico.

Where is it now?

Now introduced to British-Columbia, Washington state, Japan and Europe, where it is found on the Atlantic coast between Denmark and Spain, in Sicily and the Adriatic Sea.

How can it get here?

Contaminant of oyster spat. Hull fouling.

What you can do to prevent its arrival

Aquaculture managers and owners should avoid getting spat material from areas that are known to have slipper limpet present or nearby.

Report all sightings at the Alien Watch page of www.invasivespeciesireland.com.

Most Unwanted: Asian rapa whelk

Rapana venosa

Habitat: Marine

Threat: Threatens marine ecosystem functioning Status: Potential



Rapana venosa is a predatory marine snail which may impact both natural and cultivated populations of oysters, mussels and other molluscs. In areas where it has been introduced it has caused significant changes to the ecosystem. It has a fast growth rate and reproductive ability.

What is it?

It has a large heavy shell with a short spire. A very distinctive feature is the deep orange colour of the inside of the shell. The outer colour is variable from dull grey to red brown, with more or less conspicuous dark brown dashes on the spiral ribs, which tend to make an interrupted "vein-like" pattern throughout the entire shell.



Impact

The predatory impact of *R. venosa* has serious implications for both natural and cultivated populations of marine bivalves. *R. venosa* are very voracious predators and *Rapana* is blamed in the Black Sea for the decline of the native, edible bivalve fauna. They have caused significant changes in the ecology of bottom-dwelling organisms and have resulted in the near extinction of the Gudaut oyster. Please see the Global Invasive Species Database pages on this species for more information.

Where is it from?

Asia

Where is it now?

This species is now present in Europe and the US.

How can it get here?

Ballast water, Aquaculture and hull fouling are considered the main potential pathways to Ireland.


What you can do to prevent its arrival

Avoid hull fouling on your boats.
Never take oyster spat from an area known to have *Rapana* present and transfer to Ireland. Report all sightings at the Alien Watch page of www.invasivespeciesireland.com.

Most Unwanted: Zebra mussel

Dreissena polymorpha
Habitat: Freshwater
Threat: Fouling, negative impacts on native species and local extinction of unionids
Status: Established





The zebra mussel is perhaps one of the most well-known invasive species in Ireland. This species was first documented here in 1997. Since that date has spread to many of Irelands freshwater ecosystems and significantly altered these habitats, threatening many native species including protected and/or species of Economic value.

What is it?

The zebra mussel has a distinctive stripy shell and individuals are small, growing up to 3-4 cm in length. Unlike other freshwater mussels the zebra mussel, attaches to hard surfaces using byssal threads. Zebra mussels typically live for two to three years. They will settle on a wide range of surfaces including rocks, anchors, boat hulls, intake pipes, unionid mussels and plants.

Impact

When zebra mussels invade a waterbody there can be a wide range of economic and ecological impacts.

Zebra mussels can have varied and unpredictable ecological impacts. They can change nutrient cycles, filter out the microscopic plants called phytoplankton that form the basis of the food chain, increase water clarity and increase plant growth around lake margins. They can reduce the amount of zooplankton, an important fish food. Zebra mussels will also attach to native mussel species and this can dramatically reduce unionid populations or cause their local extinction. Changes in fish populations can also occur as zebra mussels colonize spawning grounds, change habitat and food sources. In Lough Erne, County Fermanagh, there has been an increase in water clarity and a major reduction in the amount of plankton.

Very few live specimens of the native mussel *Anodonta* are now found and the local extinction of *Anodonta* can be expected in the near future. The fish populations have also changed with much higher levels of perch recruitment.

In Northern Ireland, zebra mussels have blocked water intake pipes at Killyhevlin water works, Enniskillen, County Fermanagh and modifications were needed at a cost of over £100,000 to date.

Zebra mussels will also attach to the hulls of boats and block boat engines causing problems for boat owners. This increases maintenance costs with the need for hull cleaning or application of antifouling treatments. Zebra mussels can cause problems for anglers, with their sharp shells cutting and snagging fishing lines. Many of Northern Ireland's waterbodies support important recreational tourist fisheries that make a significant contribution to the local economy. If zebra mussels affect fish populations, this may have an economic impact.

Where are they from?

Originally from the Caspian and Black Sea region but has become invasive in Europe since the late 1700's after the construction of an extensive canal network. Zebra mussels are now present in Sweden, Finland, France, the former USSR, Germany, The Netherlands, Italy, Switzerland, Britain, Spain and North America.



Where are they in Ireland?

Zebra mussels were first discovered in Ireland in Lough Derg in 1997, but. Zebra mussels have continued to spread in Ireland and are well established in the Shannon, Boyle and Erne navigations and, to date, they have spread to over 50 lakes.

How did it get here?

The zebra mussel probably arrived in Ireland in 1994 attached to the hulls of second-hand boats imported from Britain or the Netherlands.

How are they spreading in Ireland?

The zebra mussel continues to spread by natural dispersal in connected waterbodies. The zebra mussel life cycle involves the production of millions of microscopic larvae called veligers, when water temperatures are above 12 °C (May to September in Ireland). Larvae drift in water currents and settle on hard surfaces mainly from June to October and develop the adult shape.

Zebra mussels continue to be spread by the activities of society. Hull fouling and movement of colonized material from an infested site to new waterbodies has led to this species becoming established in new lakes and rivers.

There is also a history of deliberate introduction to non impacted sites. A history of misinformation has led to some individuals believing that the zebra mussel can help 'clean up' a waterbody and remove pollution.

What you can do?

Do not introduce zebra mussels to new sites. Prevent fouling on your boat. Dry out angling equipment.

Report all sightings at the Alien Watch page of www.invasivespeciesireland.com.

Most Unwanted: Curly leaved waterweed


Lagarosiphon major

Habitat: Freshwater

Threat: Clogs waterways and negatively impacts on native species and ecosystems

Status: Established





What is it?

This species acts as an aggressive, invasive, alien plant species in Ireland but it is also regarded as a nuisance weed in its home territory. The leaves are strongly recurved and are borne in whorls of 3 or in a spiral arrangement. The long stem is brittle and easily broken (aiding dispersal). Only female plants are present and all reproduction is by fragmentation or vegetative reproduction.

Habitat

Lakes and slow flowing aquatic systems such as canals, low energy rivers and streams. The plant can grow in water up to 6m deep. In addition to occupying the full water column, plant stands produce dense vegetation on the water surface.

Impact?

Currently, in Ireland, we are at an early stage in infestation; however, significant weed stands exist in Lough, Corrib. The plant forms very dense infestations in suitable habitats and is capable of occupying the full water column in waters up to 6m deep leading to significant changes to the ecology for native plants, insects and fish.

The species is a serious threat for tourism, angling, boating and other recreational pursuits as well as conservation goals.

Where is it from?

Originating from southern Africa but the species is now invasive in mainland Europe and Britain.

Where is it now?

Present at a number of aquatic sites throughout the island.

How did it get here?

Sold as an oxygenating plant in Garden Centres and aquarium supply outlets.

How does it spread?

Spread by fragmentation via wind dispersal, boat movement, angling equipment and, possibly waterfowl. When detached stems sink they root from the nodes and establish new populations.

What you can do?

Do not grow aquatic weed species in your garden pond. Dispose of garden waste in a responsible manner. Remove all plant and animal material from equipment before leaving a water body. Report all sightings at the Alien Watch page of www.invasivespeciesireland.com.



Most Unwanted: Brown rat

Rattus norvegicus

Habitat: Terrestrial

Threat: Threatens protected species Status: Established



This species is very common throughout Ireland. It is recognized as a devastating invasive species throughout most of its global introduced range due to the impact on agriculture and on native species, sometimes causing or contributing to the extinction of species.

What is it?

The brown rat has brown fur on its back and grey fur on its belly. It is only one of two rat species in Ireland. It can be distinguished from the black rat by relatively smaller eyes and ears which are finely furred compared with the almost hairless ears of the black rat.

Impact

Both species of rats present in Ireland are known to impact on native flora and fauna. The effects these animals have maybe greater on island habitats. These species have been observed to attack and kill chicks, and adult birds, and take eggs from nest sites. Globally, these two species of rats have led to the decline, local extinction and/or total extinction of bird and other species through predation and competition. The black rat has been described to attack and kill at least 39 bird species, while the brown rat has been recorded predating on at least 53 species. Evidence of predation of small birds on islands is difficult to obtain, therefore this number is likely to be much greater. Arthropods, snails, amphibians and reptiles are generally not as well studied as bird species, and few data is available on the impacts of rats on these organisms, but again there is little doubt that the number of impacted species is very large.

Invasive rats have impacts on plant communities too. These impacts on isolated island ecosystems are extremely important and of concern to conservation managers and enthusiasts alike. Rats are omnivorous, and as well as animal species they are known to eat leaves, seeds, flowers, bark and stems of many species, some of which are endanger of extinction due to their isolation on island ecosystems and lack of adaptations to the new predator. Rats are also known to prevent the regeneration of



woodlands by damaging young saplings and adult trees.

The brown rat has out-competed the invasive black rat species on mainland Ireland.

Where is it from?

With a possible origin of North East China, this species has benefited greatly by allying itself closely to man. This species is now present in most parts of the world.

Where are they in Ireland?

Brown rats are widespread on the mainland and are present on an unknown number of offshore islands. There they reduce the reproductive success of ground nesting birds. There is still the risk of spread to other un-impacted islands.

How did it get here?

This species can stow away on freight containers and ships.

How are they spreading in Ireland?

Self-propelled. They are a highly mobile species and can swim. Stow away on both large and small vessels.

What you can do?

Prevent further spread to offshore islands by keeping boats free of rats. Use rat guards on mooring lines, where appropriate. Do not allow rats to escape off your boat and onto sensitive islands. Humanely dispatch any rat that stows away on your boat.

Report all sightings at the Alien Watch page of www.invasivespeciesireland.com.

Most Unwanted: Ship/black rat

Rattus

Habitat: Terrestrial

Threat: Threatens protected species Status: Established



The black rat is not as widespread in Ireland as the brown rat. In fact the only



known self-sustaining population is on Lambay Island off the coast of Dublin. Here this species has been protected from the Brown rat which out competes the black rat.

This species has been nominated as among 100 of the “World’s Worst” Invaders

What is it?

The black rat is much smaller than the Norway rat and can be distinguished by its ears which are larger and hairless.

Impact

Both species of rats present in Ireland are known to impact on native flora and fauna. The effects these animals have maybe greater on island habitats. These species have been observed to attack and kill chicks, and adult birds, and take eggs from nest sites. Globally, these two species of rats have led to the decline, local extinction and/or total extinction of bird and other species through predation and competition. The black rat has been described to attack and kill at least 39 bird species, while the brown rat has been recorded predating on at least 53 species. Evidence of predation of small birds on islands is difficult to obtain, therefore this number is likely to be much greater. Arthropods, snails, amphibians and reptiles are generally not as well studied as bird species, and few data is available on the impacts of rats on these organisms, but again there is little doubt that the number of impacted species is very large.

Invasive rats have impacts on plant communities too. These impacts on isolated island ecosystems are extremely important and of concern to conservation managers and enthusiasts alike. Rats are omnivorous, and as well as animal species they are known to eat leaves, seeds, flowers, bark and stems of many species, some of which are endanger of extinction due to their isolation on island ecosystems and lack of adaptations to the new Predator. Rats are also known to prevent the regeneration of woodlands by damaging young saplings and adult trees.

Where is it from?

This species of rat is native to India but now has a global distribution.

Where are they in Ireland?

The only known established and self-sustaining population of black rats is on Lambay Island off the coast of Dublin. Occasionally this species does appear around ports because of its association with shipping.

How did it get here?

This species can stow away on freight containers and ships.

How are they spreading in Ireland?



Self-propelled. They are a highly mobile species and can swim. Stow away on both large and small vessels.

What you can do?

Prevent further spread to offshore islands by keeping boats free of rats. Use rat guards where appropriate. Do not allow rats to escape off your boat and onto sensitive islands. Humanely dispatch any rat that stows away on your boat.

Report all sightings at the Alien Watch page of www.invasivespeciesireland.com.